

Obsessions and Compulsions

A self help guide

Shining a light on the future



These are the experiences of three people who suffer obsessive compulsive disorder...

“I’m afraid of catching something from other people, I fear that the germs that they carry may get on to me and I will become infected. I’m afraid I may also ‘contaminate’ my family by passing these germs on to them. I know it is silly but I feel so tense and anxious if I do touch anyone else or any surfaces – such as door handles that they have touched, that I have to come home and wash my hands many times, then wash my clothes. That makes me feel a lot better until the next contact with others. All of my own surfaces at home are washed many times each day with bleach to stop the germs. I avoid contact with other people when at all possible. Part of me realises that these fears are daft, but it’s gone on for so long now I don’t know how to stop... my family are sick of it...”

“I fear that I will harm my partner. I know that I don’t want to and I love her but thoughts often come into my head where I can picture myself harming her in some way, with a knife or by strangling her. I am so upset when I have these thoughts that I have to bring into my mind other ‘good thoughts’ such as ‘I know I love her very much’ and I say these to myself many times to get rid of the bad thoughts. I usually feel a bit better after that, until the next time the awful thoughts come into my head. I have hidden away all sharp objects and knives so that there is no risk of me doing it, and also seeing these objects brings the horrible thoughts to my mind. I spend hours each day in this mental battle... I think to myself you must be a horrible person to have those thoughts...”

“My whole day is spent checking that nothing will go wrong in the house... I can’t get out because I’m never quite sure that I’ve turned off the gas, electric appliances, water and locked the windows. No matter how often I check, my partner has to check them all for me again. I check to see if the gas fire is off, I do this five times and then can sometimes go upstairs, at other times it doesn’t feel right and I go through the whole ‘ritual’ again. If I don’t check I feel so worried I can’t bear it. I know it’s silly, but I keep thinking if something awful did happen I’d be to blame for being so careless...”

You may have had similar experiences yourself.

It is quite common for people to have such thoughts and to carry out checking actions, but if it is becoming a major part of your daily life then you may be suffering from Obsessive Compulsive Disorder. We will call it OCD in this booklet.

What is OCD?

Each person who suffers from OCD describes slightly different problems. In general people with OCD experience obsessions. These are thoughts, pictures or impulses which are usually unpleasant and come into mind when we don't want them. Many things can trigger these obsessions, and they usually leave the person feeling very anxious, uncomfortable or frightened. The compulsion is the behaviour performed in order to 'put right' the obsession. Sometimes the behaviour performed is quite irrational (and the OCD sufferer recognises this) such as counting up in sevens for seven minutes. Sometimes the behaviour is more closely related to the obsessional thought, such as washing hands many times to avoid thoughts of contamination. Most people with OCD know that their compulsions are unreasonable or 'over the top' but they feel unable to control their thoughts or change their behaviour.

Many people experience obsessions and compulsions and are able to live with this without problems. People may think about seeking help when their lives are becoming disrupted by these unwanted thoughts and actions.

What more do we know about OCD?

OCD affects us in a number of ways:

- **What we think:**
 - obsessions
 - guilty thoughts

- **How we feel:**
 - tension
 - anxiety
 - agitation

- **What we do:**
 - compulsion - (sometimes described as rituals)
 - avoidance
 - seeking reassurance

What are the symptoms of OCD?

Some of the symptoms of OCD are listed here. Most people don't experience all of these. You may want to tick any symptoms you experience regularly.

What we think – obsessions	What we do – compulsions
<ul style="list-style-type: none"> <input type="checkbox"/> Fearful thoughts or pictures in your mind about being contaminated by dangerous substances, e.g. germs, dirt, AIDS. <input type="checkbox"/> Frightening thoughts/ images that some serious harmful events will occur because of your carelessness, for example a gas explosion in the house because the cooker is left on, that the house will be burgled because of doors or windows left unlocked or that you may have knocked someone over in your car. <input type="checkbox"/> Pictures or words in your head that suggest you will harm others, especially those you care for and would never want to harm. For example that you may hurt your own child, that you may be unfaithful to your partner. <input type="checkbox"/> Pictures come into your mind of your loved ones dead. <input type="checkbox"/> Things in your life are not in the correct order or not symmetrical enough or in the right place, e.g. ornaments are out of alignment and you feel distressed by this. <input type="checkbox"/> Blasphemous or unpleasant thoughts/ pictures and doubts about your faith come into your head. 	<ul style="list-style-type: none"> <input type="checkbox"/> Check body for signs of contamination. <input type="checkbox"/> Wash/disinfect frequently. <input type="checkbox"/> Avoid going to places or touching objects that you fear may contaminate you. <input type="checkbox"/> Check feared situations/ appliances or journey route many times. <input type="checkbox"/> Avoid being the last person to leave the house. <input type="checkbox"/> Avoid responsibility. <input type="checkbox"/> Seek reassurance regularly from another person that everything is alright. <input type="checkbox"/> Avoid situations which you feel put you at risk of harming, e.g. hide kitchen knives. <input type="checkbox"/> Think something to yourself to 'put right' the frightening thoughts – neutralizing thoughts. <input type="checkbox"/> Carry out some task that will neutralize the thought, e.g. counting or saying a special word. <input type="checkbox"/> Seek reassurance from others. <input type="checkbox"/> You put things right or make them symmetrical many times until they 'feel' right. <input type="checkbox"/> You avoid contact with things that make you feel like this. <input type="checkbox"/> You pray, seek forgiveness frequently. <input type="checkbox"/> Consult religious leader/seek reassurance

How do you feel when you experience some of these obsessions?

- | | | |
|----------------------------------|--------------------------------------|------------------------------------|
| <input type="checkbox"/> fearful | <input type="checkbox"/> agitated | <input type="checkbox"/> anxious |
| <input type="checkbox"/> guilty | <input type="checkbox"/> depressed | <input type="checkbox"/> disgusted |
| <input type="checkbox"/> tense | <input type="checkbox"/> other | |

How do you feel when you have then carried out the compulsive behaviour or thoughts?


- | | | |
|--------------------------------------|---------------------------------------|---------------------------------------|
| <input type="checkbox"/> relieved | <input type="checkbox"/> cleansed | <input type="checkbox"/> calm |
| <input type="checkbox"/> relaxed | <input type="checkbox"/> less anxious | <input type="checkbox"/> disappointed |
| <input type="checkbox"/> other | | |

If you have ticked several of these thoughts, feelings and actions then you may have OCD.

Most people who have OCD find that there is a pattern in their thoughts, feelings and actions. They feel anxiety or discomfort at having the obsession and relief once they have carried out the compulsive act. This becomes a vicious cycle which strengthens itself and becomes more likely to happen again.

In addition to this the person who experiences OCD will often feel guilty and that they must be a terrible person to have such thoughts. This in turn makes the thoughts more likely to return because they are given such negative importance in the person's mind.

Research tells us that everyone has odd or distressing thoughts and pictures going through their minds at some times. Most people dismiss this from their mind as meaningless. Those who feel most guilty, distressed or disturbed by the thoughts, however, may involuntarily bring them back into their mind because of this distress. The pattern sometimes looks something like this.



Can you identify a vicious cycle that applies to your thoughts, feelings and behaviour? Try to draw it out here.

Can OCD be treated?

In the last 20 years the treatment of OCD has greatly improved and most people do make a good recovery. The most important treatments are Cognitive Behavioural Therapy (CBT), which will be described later in this booklet, and medication.

Medication

This may be prescribed by your General Practitioner or you may be offered the opportunity to see a Psychiatrist who specialises in such disorders. The medicines most commonly prescribed by doctors for OCD are antidepressant tablets, which can be very effective in the treatment of OCD even if you have no symptoms of depression. These tablets are not addictive and have few side effects. They do take a few weeks to begin to work, so if you are offered this type of treatment it will be a little time until you begin to feel the benefit.

It is important to continue with the treatment in these early weeks and to stay on the treatment as long as your doctor suggests in order to maintain full benefit.

Cognitive Behavioural Therapy

Your GP may recommend this therapy for you. This approach helps you to tackle what you think (cognition) and what to do (behaviour). Your doctor may suggest you try some of the approaches we describe in this booklet but if you require further help you may be referred to a specialist therapist. Usually this will be a Community Nurse, Nurse Therapist or a Clinical Psychologist.

What can I do to help myself?

Research has told us that the most successful way to tackle OCD is by exposure with response prevention. This literally means that you must gradually face or expose yourself to the things or situations you fear, whilst at the same time preventing yourself from carrying out your usual compulsive behaviour (checking, cleaning etc). This gradual approach means that with each stage you become less afraid of what used to trouble you and you learn by experience that no disaster occurs if you stop your compulsive behaviour.

Not all the examples in this booklet will be exactly like your experience. Use the parts that you think apply to you.

How can I make facing what I fear easier?

Exposure

We know that if we can stay in a situation where we feel anxious, gradually the anxiety will reduce – our body becomes used to the situation and we no

longer feel fearful. This is called exposure and it will help us overcome our obsessions.

For the person with OCD however, facing things we fear may seem very difficult, if not impossible.

Because of this it may be helpful to break down into smaller steps the exposure to situations or thoughts we find difficult. Begin by making a list of all situations or thoughts you find difficult. Next make an 'anxiety ladder' where those situations that you only fear a little are at the bottom and your worst feared situations are at the top. It may help to look at this example:

Washing and cleaning

Mary has a fear of being contaminated by germs which she fears she may pass on to her family. This has resulted in her restricting her lifestyle and spending many hours washing herself and disinfecting her home. She has made up the following anxiety ladder.



Mary will begin her exposure therapy at step 1 (i.e. by touching her own waste bin with gloves on) and gradually work towards step 8. She will prevent herself from frequent hand washing at these times – (see Response Prevention below). Are you able to list your own anxiety ladder?

Least feared

- 1.....
- 2.....
- 3.....
- 4.....
- 5.....
- 6.....
- 7.....
- 8.....
- 9.....
- 10.....

Most feared

Most people will at first feel greater anxiety when they begin to face the things they fear and feel an urge to perform their compulsion or ritual. The next section may help you deal with this.

How do I stop myself from carrying out the compulsive act?

It is important to break the cycle of carrying out the compulsive act or thought following exposure to the thing(s) you fear. This is called Response Prevention. There are some tips in attempting this.

1. Ask your family to help you by not offering to reassure you by checking for you or by telling you that you are not contaminated. Reassurance can stop you from confronting what you really fear.

2. Praise yourself for not carrying out the compulsion or neutralising activity. This is an important step forward.
3. Keep a note as you are carrying out the exposure therapy to show how your anxiety begins to drop. For example, touching the bin with no gloves on and without washing hands.
4. Don't substitute new compulsions for old ones. For example substituting rubbing hands continually for hand washing.
5. If stopping all compulsive behaviours at once seems impossible, try to reduce gradually the time you spend on the behaviour or the number of times it happens.

If you use this approach then gradually your anxiety will reduce.

How can I tackle negative thinking in OCD?

Sometimes people get gloomy thoughts when they have OCD especially when they begin to try and break the cycle of obsessions and compulsions by Response Prevention. Typically these thoughts are criticisms of yourself, for example, "I'm not a caring mother if I don't check things fully" or "I'm letting things get out of control, I'm a failure". These thoughts lead to low mood and unhappiness. It is important that you do not just accept these thoughts. You need to find a more balanced view. Try to:

- identify these thoughts and low mood;
- jot down the unpleasant thoughts you are having at the time;
- try and counter these thoughts by writing down arguments against them. Imagine what you would say to a friend if they had such negative thoughts about themselves;
- concentrate on and remember the good things about yourself and your life, not the bad things.

How can I tackle compulsive checking?

Mark had difficulty in leaving the house each day. He would check all appliances at least 15 times. Perhaps you have a similar problem with checking things all the time. The following example may help you understand how to tackle this kind of problem:

- write down all the things you check;
- decide which is most difficult – make an 'anxiety ladder';

- starting with step 1 on your ladder (least difficult) decide how many times you will check – try the minimum you can;
- begin with step 1 one day;
- when your anxiety about that stage is down to a low level move on to step 2.

Mark's anxiety ladder looked like this:



He began with step 1. He would check the taps only once before he left the house. At first he felt very anxious about this but gradually his fear lessened. He then moved on to step 2, the lights ... and so on.

How can I tackle obsessional thoughts where the compulsion is another thought?

Jean used to get a picture in her mind of her daughter and the words “your daughter is dead” would run into her head. She was so disturbed by this that she would ‘neutralise’ the thought in her mind by saying “she is alive and well” and would picture her daughter looking fit and healthy. She would then feel relief. This began to take up hours of Jean’s time each day and made her very unhappy. She felt she must be a terrible person. The picture seemed to get stronger and stronger. Perhaps you have a similar disturbing thought that you spend time ‘putting right’ with another thought.

The most important thing when tackling this problem is to break the cycle of having an obsessional thought and 'putting it right' with another thought, i.e. neutralising. Here are some tips!

- Don't try and get rid of the obsessional thought, just accept it. We all have odd thoughts at times. Think to yourself that it's just an odd thought, it doesn't mean anything, it doesn't mean you are a bad person.
- Do not neutralise to put the thought right – break that cycle.
- Jean tackled this by telling herself:
 - recognise that this is only a thought, it is only so upsetting if I give it too much importance. The more frightened of it I am the more it will come to mind.
 - stop trying to put the thought out of my mind – just let it fade – don't be afraid of it.
 - never 'put the thought right' by 'neutralising' (that is saying in my head "she's alive and well") – this will just strengthen the cycle.
- Remember trying not to think a thought will not help this, and can just have the opposite effect.

Test this out now - try not to think of a blue giraffe! As you can see for yourself this just brings the thought of a blue giraffe to mind! The same goes for your intrusive thoughts. Trying not to think of them may well bring them into your mind.

How can I help myself overcome OCD?

- Carefully recognise your unwanted thoughts (obsessions) and the actions you take to put them right (compulsions).
- Gradually face some of the things you fear. Work out an anxiety ladder to help you do this. Begin with the easiest step.
- Do not carry out any compulsions to reduce or neutralise your anxiety when you are facing the feared situation.
- Break the obsession compulsion cycle.
- Challenge any gloomy or critical thoughts you may have about yourself.

Where can I get further help?

- We hope you will use the exercises suggested in this booklet. They may help you overcome OCD and return to normal life.
- If you feel you are making little progress or the problem is getting worse then seek help in overcoming your problem.
- Your GP is the best person to talk to first. Your GP may suggest a talking treatment or tablets or both. They may suggest you see a mental health worker who can offer expert help with your problems.
- If you feel so distressed that you have thoughts of harming yourself, then visit your doctor as soon as possible and explain to them how you are feeling.

The following organisations and helplines may also be useful:

- **Anxiety UK**
Tel: 08444 775 774
www.anxietyuk.org.uk
Zion Community Resource Centre,
339 Stretford Road, Hulme, Manchester, M15 4ZY
Provides information and support to people suffering from anxiety disorders.
- **British Association for Counselling and Psychotherapy**
Tel: 01455 883 316
www.bacp.co.uk
BACP House, 15 St John's Business Park, Lutterworth, Leicestershire, LE17 4HB
Offers an information service providing contacts for counselling in England and Wales.
- **Carers UK**
Advice line: 0808 808 7777
www.carers.org.uk
Email: advice@carersuk.org
20 Great Dover Street, London, SE1 4LX
Provides information, advice, and support to carers and their families.
- **First Steps to Freedom**
Helpline: 0845 120 2916
Tel: 0845 841 0619
www.first-steps.org
PO Box 476, Newquay, TR7 1WQ
Provides helpline and resources for those suffering from anxiety.

- **Mental Health Matters**
 Tel: 0191 516 3500
www.mentalhealthmatters.com
 Avalon House, St Catherines Court, Sunderland Enterprise Park,
 Sunderland, SR5 3XJ
 A national organisation which provides support and information on
 employment, housing, community support and psychological services.
- **Mind Infoline**
 Tel: 0300 123 3393
www.mind.org.uk
 15-19 Broadway, Stratford, London, E15 4BQ
 Provides information on a range of topics including types of mental
 distress, where to get help, drug and alternative treatments and advocacy.
 Also provides details of help and support for people in their own area.
 Helpline available Mon - Fri, 9am - 5pm.
- **NHS Choices – Your health, your choices**
www.nhs.uk
 Information about conditions, treatments, local services and healthy lives.
- **NHS Direct**
 Tel: 0845 46 47
www.nhsdirect.nhs.uk
 Health information and advice.
- **No Panic**
 Freephone: 0808 808 0545
www.nopanic.org.uk
 93 Brands Farm Way, Telford, Shropshire, TF3 2JQ
 Support for sufferers of panic attacks, phobias, obsessive compulsive
 disorder, general anxiety disorder and tranquilliser withdrawal.
- **OCD Action**
 Tel: 0845 390 6232
www.ocdaction.org.uk
 Suite 506-507, Davina House, 137-149 Goswell Road, London, EC1V 7ET
 Provides support and information to anybody affected by OCD and works
 to raise awareness of the disorder.
- **OCD - UK**
www.ocduk.org/
 Tel: 0845 120 3778
 PO Box 8955, Nottingham, NG10 9AU
 A national support group for people with Obsessive-Compulsive Disorder.

- **PALS – Patient Advice and Liaison Service**
www.pals.nhs.uk
Provides information and advice about local NHS services and support organisations and helps sort out problems with NHS services.
- **Rethink**
Helpline: 0300 500 0927
www.rethink.org
89 Albert Embankment, London, SE1 7TP
Provides information and a helpline for anyone affected by mental health problems.
- **Samaritans**
Tel: 0845 790 9090
www.samaritans.org
PO Box 9090, Stirling, FK8 2SA
Provides confidential support for anyone in a crisis.
- **Triumph over Phobia**
Tel: 0845 600 9601
www.topuk.org
P.O. Box 3760, Bath, BA2 3WY
Provides national network of self help groups for people with phobias or OCD.

Your practice nurse or health visitor, who is based at your GP practice, will also have information about these services.

Some useful books you may like to buy or borrow from the library:

- **Living with fear**
Isaac Marks.
McGraw-Hill 2005
This self-help guide gives practical advice to people who are suffering from phobias, panic, obsessions, rituals or traumatic distress.
- **Obsessive compulsive disorder: the facts**
Padmal De Silva and Stanley Rachman
Oxford University Press 2009
In this book, the authors draw on their extensive clinical experience to give a lucid account of the nature of obsessive-compulsive behaviour. The book is intended both for those who have this disorder and for their families and friends.

- **Understanding obsessions and compulsions**

Frank Tallis

Sheldon Press

This book attempts to provide a comprehensive guide to self-help, explaining the principles of anxiety reduction, giving treatment instructions in easy-to-understand language. It covers compulsive checking, washing, hoarding, obsessional thoughts and worry, obsessional personality and depression.

- **Overcoming obsessive compulsive disorder**

David Veale and Rob Willson

Constable and Robinson 2009

With this step by step approach you can learn how to break free from the destructive cycle of obsessive behavior and regain control of your life.

- **NICE guidance**

Information for people with OCD or body dysmorphic disorder, their families and carers. Printed copies from NHS Response line

Tel: 0300 123 1002 or download from

<http://guidance.nice.org.uk/CG31/PublicInfo/pdf/English>

Support your NHS

Join our NHS Foundation Trust

Northumberland, Tyne and Wear NHS Foundation Trust provides a range of mental health, learning disability and specialist services for a large part of the North East of England and beyond. You may not think these services have very much to do with you, but mental health problems affect 1 in 4 people and there are a growing number of people with both learning and other disabilities.

As a member of our Foundation Trust you can be as active as you like, from just receiving regular information about the trust to getting involved in issues that you care about. It's free to join. Please complete and return the Membership Application Form.

Shining a light on the future



Membership Application Form

Your information

Mr Mrs Miss Other _____
(please specify)

First Name: _____ Surname: _____

Address: _____

Postcode: _____

Email: _____

How would you like us to contact you?

Post Email

About You

We are required to collect the following information about our members.

Date of Birth: _____

Male Female

Are you: White
Mixed
Asian or Asian British
Black or Black British
Other (please specify) _____

Please return your completed form to Membership Office,
Freeport NTW



Mental Health Self Help Guides

www.ntw.nhs.uk/pic

- Abuse
- Alcohol and You
- Anxiety
- Bereavement
- Controlling Anger
- Depression
- Depression and Low Mood
- Eating Disorders
- Domestic Violence
- Food for Thought
- Health Anxiety
- Hearing Voices
- Obsessions and Compulsions
- Panic
- Postnatal Depression
- Post Traumatic Stress
- Self Harm
- Shyness and Social Anxiety
- Sleep Problems
- Stress
- **Plus 3** guides for prisoners



Shining a light on the future



Written by Dr Lesley Maunder and Lorna Cameron, Consultant Clinical Psychologists, Northumberland, Tyne and Wear NHS Foundation Trust with contributions from healthcare staff and service users in Northumberland.

This information can be made available in a range of formats on request (eg Braille, audio, large print, BSL, easy read or other languages). Please contact the Patient Information Centre Tel: 0191 223 2545

Published by the Patient Information Centre

2010 Copyright, Northumberland, Tyne and Wear NHS Foundation Trust

Ref, PIC/96/0610 June 2010

www.ntw.nhs.uk/pic Tel: 0191 223 2545

Review date 2012

